

# On forcing with side conditions

David Asperó

University of East Anglia

Perspectives on Set Theory  
IMPAN, Warsaw, Nov. 2023

## $\text{MM}^{++}$ is a successful axiom (for $H(\omega_2)$ )

(1) (**Maximal forcing axiom**)  $\text{MM}^{++}$  is a consistent (relative to a supercompact), provably maximal forcing axiom relative to collections of  $\aleph_1$ -many dense sets.

(2) (**Completeness modulo forcing**) If  $\text{MM}^{++}$  holds, then  $\text{Th}(H(\omega_2)^V) = \text{Th}(H(\omega_2)^{V^P})$  for every forcing  $\mathcal{P}$  such that  $\Vdash_{\mathcal{P}} \text{MM}^{++}$  (since  $\text{MM}^{++} \Rightarrow (*)$  (A.-Schindler)).

(2) ( $\Pi_2$  **maximality**) If  $\text{MM}^{++}$  holds, then  $(H(\omega_2); \in, \text{NS}_{\omega_1}) \models \sigma$  whenever  $\sigma$  is a  $\Pi_2$  sentence such that  $(H(\omega_2); \in, \text{NS}_{\omega_1}) \models \sigma$  is forcible (again, since  $\text{MM}^{++} \Rightarrow (*)$ ); in fact, tinkering a bit with the proof that  $\text{MM}^{++} \Rightarrow (*)$  one can show that already  $\text{MM}$  is  $\Pi_2$  maximal for the theory of  $(H(\omega_2); \in)$  (A.-Schindler)).

## $MM^{++}$ is a successful axiom (for $H(\omega_2)$ )

- (1) **(Maximal forcing axiom)**  $MM^{++}$  is a consistent (relative to a supercompact), provably maximal forcing axiom relative to collections of  $\aleph_1$ -many dense sets.
- (2) **(Completeness modulo forcing)** If  $MM^{++}$  holds, then  $Th(H(\omega_2)^V) = Th(H(\omega_2)^{V^P})$  for every forcing  $\mathcal{P}$  such that  $\Vdash_{\mathcal{P}} MM^{++}$  (since  $MM^{++} \Rightarrow (*)$  (A.–Schindler)).
- (2) ( $\Pi_2$  maximality) If  $MM^{++}$  holds, then  $(H(\omega_2); \in, NS_{\omega_1}) \models \sigma$  whenever  $\sigma$  is a  $\Pi_2$  sentence such that  $(H(\omega_2); \in, NS_{\omega_1}) \models \sigma$  is forcible (again, since  $MM^{++} \Rightarrow (*)$ ); in fact, tinkering a bit with the proof that  $MM^{++} \Rightarrow (*)$  one can show that already  $MM$  is  $\Pi_2$  maximal for the theory of  $(H(\omega_2); \in)$  (A.–Schindler)).

## $MM^{++}$ is a successful axiom (for $H(\omega_2)$ )

- (1) (**Maximal forcing axiom**)  $MM^{++}$  is a consistent (relative to a supercompact), provably maximal forcing axiom relative to collections of  $\aleph_1$ -many dense sets.
- (2) (**Completeness modulo forcing**) If  $MM^{++}$  holds, then  $Th(H(\omega_2)^V) = Th(H(\omega_2)^{V^P})$  for every forcing  $\mathcal{P}$  such that  $\Vdash_{\mathcal{P}} MM^{++}$  (since  $MM^{++} \Rightarrow (*)$  (A.–Schindler)).
- (2) ( **$\Pi_2$  maximality**) If  $MM^{++}$  holds, then  $(H(\omega_2); \in, NS_{\omega_1}) \models \sigma$  whenever  $\sigma$  is a  $\Pi_2$  sentence such that  $(H(\omega_2); \in, NS_{\omega_1}) \models \sigma$  is forcible (again, since  $MM^{++} \Rightarrow (*)$ ); in fact, tinkering a bit with the proof that  $MM^{++} \Rightarrow (*)$  one can show that already  $MM$  is  $\Pi_2$  maximal for the theory of  $(H(\omega_2); \in)$  (A.–Schindler)).

Are there competitors for  $\text{MM}^{++}$  higher up? In other words, are there axioms approximating any of (1)–(3) for  $H(\omega_3)$ , or  $H(\kappa)$  for some higher  $\kappa$ ?

## $MM^{++}$ and completeness for $H(\omega_3)$

The completeness provided by (\*) for the theory of  $H(\omega_2)$  certainly doesn't extend to  $H(\omega_3)$ : Force  $\square_{\omega_1}$  by  $<\omega_2$ -distributive forcing, hence preserving (\*).

How about  $MM^{+++}$ ? Does  $MM^{+++}$  provide a complete theory, modulo forcing, for  $H(\omega_3)$ ?

The answer of course is No, but it's not so straightforward to find examples:

- (Todorčević) PFA implies  $\neg \square_{\omega_1}$ .
- (Sakai) MM implies partial square on  $S_{\omega_1}^{\omega_2}$ .
- PFA implies  $2^{\aleph_1} = \aleph_2$  (Todorčević, Veličković), so it implies  $\diamond(S_{\omega_1}^{\omega_2})$  (Shelah).
- (Baumgartner) PFA implies  $\diamond(S_{\omega_1}^{\omega_2})$ .

Given a cardinal  $\kappa$  of uncountable cofinality and a stationary set  $S \subseteq \kappa$ , *Strong Club Guessing at S*,  $\text{SCG}(S)$ , is the following statement:

There is a sequence  $(C_\delta : \delta \in S)$  such that

- for every  $\delta \in S$ ,  $C_\delta$  is a club of  $\delta$ , and
- for every club  $D \subseteq \kappa$  there are club-many  $\delta \in D$  such that if  $\delta \in S$ , then  $C_\delta \setminus \alpha \subseteq D$  for some  $\alpha < \delta$ .



## Theorem

$\text{Add}(\omega_2, \omega_3)$  forces  $\neg \text{SCG}(S)$  for every stationary  $S \subseteq S_\omega^{\omega_2}$ .  
Hence, if  $\text{MM}^{++}$  holds, then forcing with  $\text{Add}(\omega_2, \omega_3)$  yields a model of  $\text{MM}^{++} + \neg \text{SCG}(S)$  for every stationary  $S \subseteq S_\omega^{\omega_2}$ .

## Theorem

Let  $\kappa$  be a supercompact cardinal, and let  $\mathcal{P}$  be the standard RCS-iteration of length  $\kappa$  forcing  $\text{MM}^{++}$ . Let  $S = (S_\omega^{\omega_2})^V$ . Then  $\mathcal{P} * \dot{Q}(S)$  forces  $\text{MM}^{++} + \text{SCG}(S)$ . Here,  $\dot{Q}(S)$  is a natural  $\aleph_1$ -support iteration of length  $\omega_3$  for adding some  $(\dot{C}_\delta : \delta \in S)$  and then shooting clubs through

$$\{\delta \in \omega_2 : \delta \in S \Rightarrow \dot{C}_\delta \setminus \alpha \subseteq \dot{D}_\alpha \text{ for some } \alpha < \delta\},$$

where  $\dot{D}_\alpha$  is a club of  $\omega_2$ .

**Question:** Is there any forcible  $\Sigma_2$  axiom  $A$  deciding the theory of  $H(\omega_3)$  modulo forcing?

## Theorem

$\text{Add}(\omega_2, \omega_3)$  forces  $\neg \text{SCG}(S)$  for every stationary  $S \subseteq S_\omega^{\omega_2}$ .  
Hence, if  $\text{MM}^{++}$  holds, then forcing with  $\text{Add}(\omega_2, \omega_3)$  yields a model of  $\text{MM}^{++} + \neg \text{SCG}(S)$  for every stationary  $S \subseteq S_\omega^{\omega_2}$ .

## Theorem

Let  $\kappa$  be a supercompact cardinal, and let  $\mathcal{P}$  be the standard RCS-iteration of length  $\kappa$  forcing  $\text{MM}^{++}$ . Let  $S = (S_\omega^{\omega_2})^V$ . Then  $\mathcal{P} * \dot{Q}(S)$  forces  $\text{MM}^{++} + \text{SCG}(S)$ . Here,  $\dot{Q}(S)$  is a natural  $\aleph_1$ -support iteration of length  $\omega_3$  for adding some  $(\dot{C}_\delta : \delta \in S)$  and then shooting clubs through

$$\{\delta \in \omega_2 : \delta \in S \Rightarrow \dot{C}_\delta \setminus \alpha \subseteq \dot{D}_\alpha \text{ for some } \alpha < \delta\},$$

where  $\dot{D}_\alpha$  is a club of  $\omega_2$ .

**Question:** Is there any forcible  $\Sigma_2$  axiom  $A$  deciding the theory of  $H(\omega_3)$  modulo forcing?

# Limitations on completeness

## Theorem

(Woodin) Suppose the  $\Omega$  conjecture and the  $AD^+$ -conjecture are true in all set-generic extensions. Then there is no forcible  $\Sigma_2$  axiom  $A$  such that  $A$  provides, modulo forcing, a complete theory for  $\Sigma_3^2$  sentences.

## Theorem

(Woodin) Suppose the  $\Omega$  conjecture holds and there is a proper class of Woodin cardinal. Then there is no forcible  $\Sigma_2$  axiom  $A$  such that  $A$  provides, modulo forcing, a complete theory for  $H(\delta_0^+)$ , where  $\delta_0$  is the first Woodin cardinal.

# Limitations on completeness

## Theorem

(Woodin) Suppose the  $\Omega$  conjecture and the  $AD^+$ -conjecture are true in all set-generic extensions. Then there is no forcible  $\Sigma_2$  axiom  $A$  such that  $A$  provides, modulo forcing, a complete theory for  $\Sigma_3^2$  sentences.

## Theorem

(Woodin) Suppose the  $\Omega$  conjecture holds and there is a proper class of Woodin cardinal. Then there is no forcible  $\Sigma_2$  axiom  $A$  such that  $A$  provides, modulo forcing, a complete theory for  $H(\delta_0^+)$ , where  $\delta_0$  is the first Woodin cardinal.

## High $\Pi_2$ maximality?

$\Pi_2$  forcing maximality for the theory  $H(\omega_3)$  is false, at least in the presence of a Mahlo cardinal:

Both  $\square_{\omega_1}$  and  $\neg\square_{\omega_1}$  can be forced, and  $\square_{\omega_1}$  is  $\Sigma_1(\omega_2)$  over  $H(\omega_3)$ .

**Question:** Does ZFC prove that  $\Pi_2$  forcing maximality for the theory  $H(\omega_3)$  is false? Does it in fact prove that there is a  $\Sigma_1(\omega_2)$  sentence  $\sigma$  such that both  $\sigma$  and  $\neg\sigma$  are forcible?

A vague question:

**Question:** Can there (still) be any reasonable successful analogue of  $MM^{++}$ , as forcing axiom, for  $H(\omega_3)$  or higher up?

- Such an analogue of  $MM^{++}$ , if it extends  $FA_{\omega_2}(\{\text{Cohen}\})$ , should presumably imply  $2^{\aleph_0} = \aleph_3$ .
- Alternatively, we could instead focus, in the context of CH, on interesting classes  $\Gamma$  of countably closed forcings.

## Strong properness

(Mitchell) A partial order  $\mathcal{P}$  is *strongly proper* iff for every large enough cardinal  $\theta$ , every countable  $M \preceq H(\theta)$  such that  $\mathcal{P} \in M$ , and every  $p \in \mathcal{P} \cap M$  there is some  $q \leq_{\mathcal{P}} p$  which is *strongly  $(M, \mathcal{P})$ -generic*, i.e., for every  $q' \leq_{\mathcal{P}} q$  there is some  $\pi_M(q') \in \mathcal{P} \cap M$  such that every  $r \in \mathcal{P} \cap M$  with  $r \leq_{\mathcal{P}} \pi_M(q')$  is compatible with  $q'$ .

Examples of strongly proper posets:

- Cohen forcing
- Baumgartner's forcing for adding a club of  $\omega_1$  with finite conditions.

**Caution:** ccc does not imply strongly proper. In fact, most ccc forcings are not strongly proper.



## Some basic facts

### Fact

*If  $\mathcal{P}$  is strongly proper,  $M \preceq H(\theta)$  is countable,  $\mathcal{P} \in M$ ,  $q$  is strongly  $(M, \mathcal{P})$ -generic,  $G \subseteq \mathcal{P}$  is generic over  $V$ , and  $q \in G$ , then  $G \cap M$  is  $\mathcal{P} \cap M$ -generic over  $V$ .*

### Corollary

*Every  $\omega$ -sequence of ordinals added by a strongly proper forcing notion is in a generic extension of  $V$  by Cohen forcing.*

### Lemma

*(Neeman) Suppose  $\mathcal{P}$  is strongly proper. Then  $\mathcal{P}$  does not add new  $ht(T)$ -branches through trees  $T$  such that  $cf(ht(T)) \geq \omega_1$ .*

## Some pure side condition forcings (chains)

(1) (Todorćević)  $\mathbb{C}_1$ : conditions are chains  $\mathcal{C} = \{M_0, \dots, M_n\}$  with  $M_i \preceq H(\theta)$ ,  $|M_i| = \aleph_0$ ,  $M_i \in M_{i+1}$  for all  $i$ .

- $\mathbb{C}_1$  is strongly proper for countable models.
- $\mathbb{C}_1$  covers  $H(\theta)^V$  by an  $\in$ -chain of length  $\omega_1$  of countable models in  $V$ .

(2) (Neeman)  $\mathbb{C}_2$ : conditions are  $\mathcal{C} = \{Q_0, \dots, Q_n\}$ , where

- (a)  $Q_i$  is either a countable  $M \preceq (\theta)$  or  $N \preceq (\theta)$  such that  $|N| = \aleph_1$  and  $N$  internally club (IC).
- (b)  $Q_i \in Q_{i+1}$  for all  $i < n$ .
- (c) If  $N, M \in \mathcal{N}$ ,  $N \in M$ ,  $|N| = \aleph_1$ ,  $|M| = \aleph_0$ , then  $N \cap M \in \mathcal{C}$ .

- $\mathbb{C}_2$  is strongly proper for countable models and IC models of size  $\aleph_1$ .
- $\mathbb{C}_2$  covers  $H(\theta)^V$  by an  $\in$ -chain of length  $\omega_1$  of  $\aleph_1$ -sized models in  $V$ .

## A limitation

### Fact

(Veličković) *The natural pure side condition forcing  $\mathbb{C}_3$  for three types of models (say countable, size  $\aleph_1$  IC, and size  $\aleph_2$  IC) doesn't work.*

## More pure side condition forcings (symmetric systems)

- (3) (Todorčević, A.–Mota, ...)  $\mathbb{S}_1$ : conditions are finite collections  $\mathcal{N}$  of countable  $M \preceq H(\theta)$  such that
- (a) For all  $M_0, M_1 \in \mathcal{N}$ , if  $\delta_{M_0} = \delta_{M_1}$  ( $\delta_M = M \cap \omega_1$ ), then  $M_0 \cong M_1$  and the isomorphism

$$\Psi_{M_0, M_1} : M_0 \longrightarrow M_1$$

is the identity on  $M_0 \cap M_1$ .

- (b) For all  $M_0, M_1 \in \mathcal{N}$ , if  $\delta_{M_0} = \delta_{M_1}$ , then  $\Psi_{M_0, M_1} \text{ “ } \mathcal{N} \cap M_0 = \mathcal{N} \cap M_1 \text{ ”}$ .
- (c) For all  $M_0, M_1 \in \mathcal{N}$ , if  $\delta_{M_0} < \delta_{M_1}$ , then there is some  $M'_1 \in \mathcal{N}$  such that  $M_0 \in M'_1$  and  $\delta_{M'_1} = \delta_{M_1}$ .

( $\mathcal{N}$  is a *symmetric system*)

- $\mathbb{S}_1$  is strongly proper for countable models.
- (CH)  $\mathbb{S}_1$  has the  $\aleph_2$ -c.c. and preserves CH.

(4) (Gallart, Hoseini Naveh)  $\mathbb{S}_2$ : conditions are *symmetric systems*  $\mathcal{N}$  of models of two types (countable and IC of size  $\aleph_1$ ).

- (a) Natural combination of Neeman's notion of two-type chain of models ( $\mathbb{C}_2$ ) and the notion of symmetric system ( $\mathbb{S}_1$ ).
- (b) Given two models  $M_0, M_1 \in \mathcal{N}$  of the same height  $\epsilon_M$  ( $= \sup(M \cap \omega_2)$ ), we ask that in fact

$$(\text{Hull}(M_0, \omega_1); \in, M_0) \cong (\text{Hull}(M_1, \omega_1); \in, M_1)$$

- $\mathbb{S}_2$  is strongly proper for countable models and for  $\aleph_1$ -sized IC models.
- ( $2^{\aleph_1} = \aleph_2$ )  $\mathbb{S}_2$  has the  $\aleph_3$ -c.c. and preserves  $2^{\aleph_1} = \aleph_2$ .

## An application of $\mathbb{S}_2$

A *strong*  $\omega_3$ -chain of subsets of  $\omega_1$  is a sequence  $(X_i : i < \omega_3)$  of subsets of  $\omega_1$  such that for all  $i_0 < i_1$ ,

- $X_{i_0} \setminus X_{i_1}$  is finite and
- $|X_{i_1} \setminus X_{i_0}| = \aleph_1$ .

**Theorem\*** (A.–Gallart) (GCH) There is a forcing notion  $\mathcal{P}$  with the following properties.

- (1)  $\mathcal{P}$  is proper for countable models and for IC models of size  $\aleph_1$ .
- (2)  $\mathcal{P}$  has the  $\aleph_3$ -chain condition.
- (3)  $\mathcal{P}$  forces the existence of a strong  $\omega_3$ -chain of subsets of  $\omega_1$ .

$\mathcal{P}$  uses side conditions from  $\mathbb{S}_2$  in a crucial way.

This result is optimal:

## Theorem

(Inamdar) *There is no strong  $\omega_3$ -chain of subsets of  $\omega_2$ .*

A *strong  $\omega_3$ -chain of functions from  $\omega_1$  into  $\omega_1$*  is a sequence  $(h_i : i < \omega_3)$  of functions  $h_i : \omega_1 \rightarrow \omega_1$  such that for all  $i_0 < i_1 < \omega_3$ ,

$$\{\tau \in \omega_1 : h_{i_1}(\tau) \leq h_{i_0}(\tau)\}$$

is finite.

**Question:** Is it consistent to have a strong  $\omega_3$ -chain of functions from  $\omega_1$  into  $\omega_1$ ?

## Extending strong properness to $\kappa > \omega$

The notion of strong properness can be naturally extended to higher cardinals:

Suppose  $\kappa$  is an infinite regular cardinal such that  $\kappa^{<\kappa} = \kappa$ . A partial order  $\mathcal{P}$  is  $\kappa$ -strongly proper iff for every  $M \preccurlyeq H(\theta)$  such that  $\mathcal{P} \in M$  and such that

- $|M| = \kappa$ , and
- ${}^{<\kappa}M \subseteq M$ ,

every  $\mathcal{P}$ -condition in  $M$  can be extended to a strongly  $(M, \mathcal{P})$ -generic condition.



We will need the following closure property:

Given an infinite regular cardinal  $\kappa$ , a partial order  $\mathcal{P}$  is  $<\kappa$ -directed closed with greatest lower bounds in case every directed subset  $X$  of  $\mathcal{P}$  (i.e., every finite subset of  $X$  has a lower bound in  $\mathcal{P}$ ) such that  $|X| < \kappa$  has a greatest lower bound in  $\mathcal{P}$ .

We will also say that  $\mathcal{P}$  is  $\kappa$ -lattice.

All facts about strongly proper (i.e.,  $\omega$ -strongly proper) forcing we have seen extend naturally to  $\kappa$ -strongly proper forcing notion which are  $\kappa$ -lattice (assuming  $\kappa^{<\kappa} = \kappa$ ).

For example, every  $\kappa$ -sequence of ordinals added by a forcing in this class belongs to a generic extension by adding a Cohen subset of  $\kappa$ .

## Lemma

(Reflection Lemma) Let  $\kappa$  be an infinite regular cardinal such that  $\kappa^{<\kappa} = \kappa$ . Suppose  $\mathcal{P}$  is a  $\kappa$ -lattice and  $\kappa$ -strongly proper forcing. If  $\theta$  is large enough and  $(M_i)_{i < \kappa^+}$  is a  $\subseteq$ -continuous  $\varepsilon$ -chain of elementary submodels of  $H(\theta)$  such that  $\mathcal{P} \in M_i$ ,  $|M_i| = \kappa$ , and  ${}^{<\kappa}M_i \subseteq M_i$  for all  $i \in \mathcal{S}_\kappa^+$ , then  $\mathcal{P} \cap N$  is  $\kappa$ -lattice and  $\kappa$ -strongly proper, for  $N = \bigcup_{i < \kappa^+} M_i$ .

## Proof.

Let  $\chi$  large enough and  $M^* \preceq H(\chi)$  such that  $\mathcal{P}, (M_i)_{i < \kappa^+} \in M^*$ ,  $|M^*| = \kappa$  and  ${}^{<\kappa}M^* \subseteq M^*$ . Then  $M^* \cap N = M_\delta \in N$  for  $\delta = M^* \cap \kappa^+$ . But every strongly  $(M_\delta, \mathcal{P})$ -generic is strongly  $(M^*, \mathcal{P} \cap N)$ -generic. □

Compare the above reflection property with the reflection of  $\kappa$ -c.c. forcing to substructures  $M$  such that  ${}^{<\kappa}M \subseteq M$ .

## Theorem

(A.–Cox–Karagila–Weiss) Assume GCH, and let  $\kappa$  be infinite regular cardinal. Then there is a  $\kappa$ -lattice and  $\kappa$ -strongly proper forcing  $\mathcal{P}$  which forces  $2^\kappa = \kappa^{++}$  together with the  $\kappa$ -Str PFA (=  $FA_{\kappa^+}(\kappa\text{-lattice} + \kappa\text{-strongly proper})$ ).

**Proof sketch:** Let  $\theta = \kappa^{++}$ . By first forcing with  $\text{Coll}(\kappa^+, <\theta)$ , we may assume that  $\diamond(S_{\kappa^+}^\theta)$  holds.

Our forcing  $\mathcal{P}$  is  $\mathcal{P}_\theta$ , where  $(\mathcal{P}_\alpha, \dot{Q}_\beta, : \alpha \in E \cup \{\theta\}, \beta \in E)$ ,  $E \subseteq S_{\kappa^{++}}^\theta$ , is a  $<\kappa$ -support iteration à la Neeman with side conditions from  $\mathbb{C}_2(\mathcal{S}, \mathcal{T})$ , for

$$\mathcal{S} = \{M : |M| = \kappa, <^\kappa M \subseteq M\}$$

and

$$\mathcal{T} = \{N_\alpha : \alpha \in E\},$$

where  $(N_\alpha : \alpha \in E)$  is some filtration of  $H(\theta)$ .

Condition are  $p = (w_p, C_p)$ , where

- $\text{dom}(w_p) \in [\theta]^{<\kappa}$ ;
- $C_p \in \mathbb{C}_2(\mathcal{S}, \mathcal{T})$ ;
- for all  $\alpha \in \text{dom}(w_p)$ ,  $N_\alpha \in C_p$  and

$(w_p \upharpoonright \alpha, \mathcal{N}_p \cap N_\alpha) \Vdash_{\mathcal{P}_\alpha}$  “ $w_p(\alpha)$  is strongly  $(M[\dot{G}_\alpha], \dot{Q}_\alpha)$ -generic”

for all  $M \in C_p \cap \mathcal{S}$  with  $\alpha \in M$ .

At stage  $\alpha$ , if our diamond feeds us a  $\mathcal{P}_\alpha$ -name  $\dot{Q}_\alpha$  for a  $\kappa$ -lattice  $\kappa$ -strongly proper forcing, then we let  $\dot{Q}_\alpha = \dot{Q}_\alpha$ .

The Reflection Property is used to show that our construction captures  $\kappa$ -strongly proper forcings of arbitrary size.

The proof uses the fact that every  $\kappa$ -sequence of ordinals is in a  $\kappa$ -Cohen extension since each  $\mathcal{P}_\alpha$  is  $\kappa$ -lattice and  $\kappa$ -strongly proper, which enables a typical model  $N_\alpha \in \mathcal{T}$  to have access to the relevant  $\mathcal{P}_\alpha$ -names for  $\kappa$ -sized elementary submodels  $M$  (so the relevant  $\dot{Q}_\alpha$ 's are in fact such that  $\Vdash_{\mathcal{P}_\alpha} \dot{Q}_\alpha$  is  $\kappa$ -strongly proper).

Also: The proof crucially uses the fact that our forcings are  $\kappa$ -lattice (it would not work if we just assumed  $<\kappa$ -directed closedness).  $\square$

$\kappa$ -Str PFA does not decide  $2^\kappa$ . In fact:

### Theorem

Assume GCH, and let  $\kappa < \kappa^+ < \kappa^{++} \leq \theta$  be infinite regular cardinals. Suppose  $\diamond(S_{\kappa^+}^{\kappa^{++}})$  holds. Then there is a  $\kappa$ -lattice and  $\kappa$ -strongly proper forcing  $\mathcal{P}$  which forces  $2^\kappa = \theta$  together with  $\kappa$ -Str PFA.

**Proof sketch:** We build an iteration

$$(\mathcal{P}_\alpha, \dot{Q}_\beta : \alpha \in E \cup \{\kappa^{++}\}, \beta \in E)$$

as before, except that at each stage  $\alpha \in E$  now we look at whether our diamond feeds us a  $\mathcal{P}_\alpha \times \text{Add}(\kappa, \kappa^+)$ -name  $\dot{Q}_\alpha$  for a  $\kappa$ -lattice and  $\kappa$ -strongly proper poset. If so we let  $\dot{Q}_\alpha = \text{Add}(\kappa, \kappa^+) * \dot{Q}_\alpha$ .

The forcing witnessing the theorem is

$$\mathcal{P} = \mathcal{P}_{\kappa^{++}} \times \text{Add}(\kappa, \theta)$$

To see this, take a  $\kappa$ -lattice  $\kappa$ -strongly proper forcing in the extension via  $\mathcal{P}$ . By the Reflection Property it reflects to a forcing of size  $\kappa^+$ . Let  $\dot{Q}$  be a  $\mathcal{P}$ -name for the corresponding forcing.

By  $\kappa^{++}$ -c.c. of  $\mathcal{P}$  we may identify  $\dot{Q}$  with a  $\mathcal{P}_{\kappa^{++}} \times \text{Add}(\kappa, \kappa^+)$ -name, which we may code by a subset of  $\kappa^{++}$ . Now we use our diamond to capture  $\dot{Q}$  as in the proof of the previous theorem.



Again, we use the fact that every  $\kappa$ -sequence of ordinals in the final model is in a  $\kappa$ -Cohen extension since  $\mathcal{P}_\alpha \times \text{Add}(\kappa, \kappa^+)$  is  $\kappa$ -lattice and  $\kappa$ -strongly proper.



As far as I know this is the first example of a forcing axiom  $\text{FA}_{\kappa^+}(\Gamma)$  such that  $\text{FA}_{\kappa^{++}}(\Gamma)$  is false but nevertheless  $\text{FA}_{\kappa^+}(\Gamma)$  is compatible with  $2^\kappa$  arbitrarily large:

To see that  $\text{FA}_{\kappa^{++}}(\kappa\text{-lattice} + \kappa\text{-strongly proper})$  is false, look at the forcing  $\mathbb{P}$  of  $<_\kappa$ -length  $\in$ -chains of suitable models  $N \preceq H(\kappa^{++})$  of size  $\kappa$  (this is  $\mathbb{C}_1$  in this context). An application of  $\text{FA}_{\kappa^{++}}(\{\mathbb{P}\})$  would cover  $\kappa^{++}$  with a  $\kappa^+$ -chain of models of size  $\kappa$ .

Again, we use the fact that every  $\kappa$ -sequence of ordinals in the final model is in a  $\kappa$ -Cohen extension since  $\mathcal{P}_\alpha \times \text{Add}(\kappa, \kappa^+)$  is  $\kappa$ -lattice and  $\kappa$ -strongly proper.



As far as I know this is the first example of a forcing axiom  $\text{FA}_{\kappa^+}(\Gamma)$  such that  $\text{FA}_{\kappa^{++}}(\Gamma)$  is false but nevertheless  $\text{FA}_{\kappa^+}(\Gamma)$  is compatible with  $2^\kappa$  arbitrarily large:

To see that  $\text{FA}_{\kappa^{++}}(\kappa\text{-lattice} + \kappa\text{-strongly proper})$  is false, look at the forcing  $\mathbb{P}$  of  $<\kappa$ -length  $\in$ -chains of suitable models  $N \preceq H(\kappa^{++})$  of size  $\kappa$  (this is  $\mathbb{C}_1$  in this context). An application of  $\text{FA}_{\kappa^{++}}(\{\mathbb{P}\})$  would cover  $\kappa^{++}$  with a  $\kappa^+$ -chain of models of size  $\kappa$ .

# Applications of $\kappa$ -Str PFA

Not many.

- $\mathfrak{d}(\kappa) > \kappa^+$
- The covering number of natural meagre ideals is  $> \kappa^+$ .
- Weak failures of Club-Guessing at  $\kappa$ .

## Relaxing strongness?

Let us say that a forcing  $\mathcal{P}$  is  $\kappa$ -MRP-*strongly proper* if for every large enough  $\theta$ , every  $M \preccurlyeq H(\theta)$  of size  $\kappa$  such that  ${}^{<\kappa}M \subseteq M$  and  $\mathcal{P} \in M$ , and every  $p \in M \cap \mathcal{P}$  there is  $q \leq_{\mathcal{P}} p$  such that for every  $q' \leq_{\mathcal{P}} q$ ,

$$\mathcal{X}_{q'} = \{X \in [M]^{\kappa} : \exists \pi_X(q') \in \mathcal{P} \cap X \forall r \leq_{\mathcal{P}} \pi_X(q'), r \in X \longrightarrow r \Vdash_{\mathcal{P}} q'\}$$

is  $M$ -stationary (i.e., for every club  $E \in M$  there is some  $X \in E \cap \mathcal{X}_{q'} \cap M$ ).

$\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice and } \kappa\text{-MRP-strongly proper}\})$  implies a natural high analogue of MRP which in turn implies  $2^{\kappa^+} = \kappa^{++}$ .

### Theorem

*Suppose  $\kappa \geq \omega_1$  is a regular cardinal and  $\kappa^{<\kappa} = \kappa$ . Then*

$$\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice, } \kappa^+\text{-c.c., and } \kappa\text{-MRP-strongly proper}\})$$

*is false.*

**Proof sketch:** For the proof we use...

## Relaxing strongness?

Let us say that a forcing  $\mathcal{P}$  is  $\kappa$ -MRP-*strongly proper* if for every large enough  $\theta$ , every  $M \preccurlyeq H(\theta)$  of size  $\kappa$  such that  ${}^{<\kappa}M \subseteq M$  and  $\mathcal{P} \in M$ , and every  $p \in M \cap \mathcal{P}$  there is  $q \leq_{\mathcal{P}} p$  such that for every  $q' \leq_{\mathcal{P}} q$ ,

$$\mathcal{X}_{q'} = \{X \in [M]^{\kappa} : \exists \pi_X(q') \in \mathcal{P} \cap X \forall r \leq_{\mathcal{P}} \pi_X(q'), r \in X \longrightarrow r \Vdash_{\mathcal{P}} q'\}$$

is  $M$ -stationary (i.e., for every club  $E \in M$  there is some  $X \in E \cap \mathcal{X}_{q'} \cap M$ ).

$\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice and } \kappa\text{-MRP-strongly proper}\})$  implies a natural high analogue of MRP which in turn implies  $2^{\kappa^+} = \kappa^{++}$ .

### Theorem

Suppose  $\kappa \geq \omega_1$  is a regular cardinal and  $\kappa^{<\kappa} = \kappa$ . Then

$$\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice, } \kappa^+\text{-c.c., and } \kappa\text{-MRP-strongly proper}\})$$

is false.

**Proof sketch:** For the proof we use...

# An inconsistent uniformization principle

## Theorem

(Shelah) Let  $\kappa \geq \omega_1$  be a regular cardinal and let  $\langle C_\alpha : \alpha \in S_\kappa^{\kappa^+} \rangle$  be a club-sequence. Then there is a sequence

$$\langle f_\alpha : \alpha \in S_\kappa^{\kappa^+} \rangle$$

of colourings, with  $f_\alpha : C_\alpha \rightarrow \{0, 1\}$  for all  $\alpha$ , for which there is no function

$$G : \kappa^+ \rightarrow 2$$

such that for all  $\alpha \in S_\kappa^{\kappa^+}$ ,

$$G(\xi) = f_\alpha(\xi)$$

for club-many  $\xi \in C_\alpha$ .

Now let  $\langle \mathcal{C}_\alpha : \alpha \in \mathcal{S}_\kappa^{\kappa^+} \rangle$  be a club-sequence and  $\langle f_\alpha : \alpha \in \mathcal{S}_\kappa^{\kappa^+} \rangle$  be a sequence of colourings which cannot be club-uniformized.

Let  $\mathcal{P}$  be the forcing consisting of  $<\kappa$ -sized functions  $p$  with  $\text{dom}(p) \subseteq \mathcal{S}_\kappa^{\kappa^+}$  such that

- (1) for all  $\alpha \in \text{dom}(p)$ ,  $p(\alpha) < \alpha$ , and
- (2) for all  $\alpha_0 < \alpha_1$  in  $\text{dom}(p)$ , if  $\xi \in (\mathcal{C}_{\alpha_0} \setminus p(\alpha_0)) \cap (\mathcal{C}_{\alpha_1} \setminus p(\alpha_1))$ , then  $f_{\alpha_0}(\xi) = f_{\alpha_1}(\xi)$ .

Then  $\mathcal{P}$  is  $\kappa^+$ -c.c.,  $\kappa$ -lattice, and  $\kappa$ -MRP-strongly proper, so an application of  $\text{FA}_{\kappa^+}(\{\mathcal{P}\})$  gives us a function  $G : \kappa^+ \rightarrow \{0, 1\}$  which in fact uniformizes  $\langle f_\alpha : \alpha \in \mathcal{S}_\kappa^{\kappa^+} \rangle$  modulo co-bounded sets — for each  $\alpha \in \mathcal{S}_\kappa^{\kappa^+}$  there is  $p(\alpha) < \alpha$  such that  $G(\xi) = f_\alpha(\xi)$  for all  $\xi \in \mathcal{C}_\alpha \setminus p(\alpha)$ .  $\square$

## Getting rid of g.l.b.'s?

No:

### Theorem

(Shelah) Suppose  $\kappa \geq \omega_1$  is a regular cardinal and  $\kappa^{<\kappa} = \kappa$ .  
Then

$FA_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } <_{\kappa}\text{-directed closed, } \kappa^+\text{-c.c., and } \kappa\text{-strongly proper}\})$

is false.

### Proof.

Similar as previous proof, with a natural forcing for adding

$G : \kappa^+ \rightarrow \{0, 1\}$  and clubs  $D_\alpha \subseteq C_\alpha$  (for  $\alpha \in S_{\kappa^+}^{\kappa^+}$ ) such that  
 $G(\xi) = f_\alpha(\xi)$  for all  $\alpha$  and all  $\xi \in D_\alpha$ . □



## $\kappa$ -strong semiproperness

Let  $\kappa$  be an infinite regular cardinal such that  $\kappa^{<\kappa} = \kappa$ . Let us say that a forcing notion  $\mathcal{P}$  is  $\kappa$ -strongly semiproper if and only if for every large enough  $\theta$  and every  $M \preceq H(\theta)$  such that  $\mathcal{P} \in M$ ,  $|M| = \kappa$ , and  ${}^{<\kappa}M \subseteq M$ , every  $p \in \mathcal{P} \cap M$  can be extended to some  $q \in \mathcal{P}$  which is  $\kappa$ -strongly  $(M, \mathcal{P})$ -semigeneric, i.e., there is some  $\sigma \in [H(\theta)]^{\leq \kappa}$  such that

- (1)  $\text{Hull}(M, \sigma) \cap \kappa^+ = M \cap \kappa^+$ , and
- (2)  $q$  is strongly  $(\text{Hull}(M, \sigma), \mathcal{P})$ -generic.

Given infinite regular  $\kappa$ , let the  $\kappa$ -Strongly Semiproper Forcing Axiom be

$$\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice and } \kappa\text{-strongly semiproper}\})$$

# A family of reflection principles

Given an infinite regular  $\kappa$  and a cardinal  $\mu \leq \kappa$ , let

$$\text{SRP}(\kappa^+, \mu)$$

be the following reflection principle: Suppose  $X$  is a set and  $\mathcal{S} \subseteq [X]^\kappa$ . If  $\lambda$  is such that  $X \in H(\lambda)$ , there is a  $\subseteq$ -continuous  $\in$ -chain  $(M_i)_{i < \kappa^+}$  such that for each  $i < \kappa^+$ ,  $M_i \preceq H(\lambda)$  and  $|M_i| = \kappa$ , and if  $\text{cf}(i) = \kappa$ :

- $M_i \cap X \notin \mathcal{S}$  if and only if there is no  $\sigma \in [X]^{\leq \mu}$  such that
  - (a)  $\text{Hull}(M_i \cup \sigma)$  is a  $\kappa^+$ -end-extension of  $M$  (i.e.,  $\text{Hull}(M_i \cup \sigma) \cap \kappa^+ = M_i \cap \kappa^+$ ), and
  - (b)  $\text{Hull}(M_i \cup \sigma) \cap X \in \mathcal{S}$ .

Easy: The  $\kappa$ -Strongly Semiproper Forcing Axiom implies  $\text{SRP}(\kappa^+, \kappa)$ .

## Theorem

For every  $\kappa \geq \omega_1$ ,  $\text{SRP}(\kappa^+, \omega)$  is false. In particular, the  $\kappa$ -Strongly Semiproper Forcing Axiom is false.

**Proof:** Let  $\mathcal{S}$  be the collection of  $X \in [\kappa^{++}]^\kappa$  such that  $\text{cf}(X) = \omega$ .

By an application of  $\text{SRP}(\kappa^+, \omega)$  to  $\mathcal{S}$  there is a  $\subseteq$ -continuous  $\in$ -chain  $(M_i)_{i < \kappa^+}$  of models of size  $\kappa$  such that for each  $i < \kappa^+$  such that  $\text{cf}(i) = \kappa$ , if

$$\text{cf}(M_i \cap \kappa^{++}) \neq \omega,$$

then there is no countable  $\sigma \subseteq \kappa^{++}$  such that

- $\text{Hull}(M_i \cup \sigma) \cap \kappa^+ = M_i \cap \kappa^+$  and
- $\text{cf}(\text{Hull}(M_i \cup \sigma) \cap \kappa^{++}) = \omega$ .

## Claim:

$S = \{i \in S_{\kappa}^{\kappa^+} : \text{there is no countable } \sigma \subseteq \kappa^{++} \text{ as above for } M_i\}$

cannot be stationary: Suppose  $S$  is stationary. Let  $\alpha \in \kappa^{++}$ ,  $\text{cf}(\alpha) = \omega$ , such that  $F''[\alpha]^{<\omega} \cap \kappa^{++} \subseteq \alpha$  for some  $F : [H(\lambda)]^{<\omega} \rightarrow H(\lambda)$  generating club of elementary submodels  $R$  such that  $(M_i)_{i < \kappa^+} \in R$ .

Now we can easily find  $X \subseteq \alpha$  cofinal in  $\alpha$ , such that  $R = F''[X]^{<\omega}$  is such that  $|R| = \kappa$  and  $i := R \cap \kappa^+ \in S$ . Let  $\sigma \subseteq X$  be countable and cofinal in  $X$ . But then  $R$  is a  $\kappa^+$ -end-extension of  $M_i$  and  $\text{cf}(R \cap \kappa^{++}) = \omega$ , and so  $\sigma$  witnesses that  $M_i \notin S$ . Contradiction.  $\square$

Now we get club-many  $i$  such that if  $\text{cf}(i) = \kappa$ , then  $\text{cf}(M_i \cap \kappa^{++}) = \omega$ . But this is impossible since  $(\sup(M_i \cap \kappa^{++})) : i < \kappa^+$  is strictly increasing and continuous and therefore  $\text{cf}(M_i \cap \kappa^{++}) = \kappa > \omega$  if  $\text{cf}(i) = \kappa$ .  $\square$

## Claim:

$S = \{i \in S_{\kappa}^{\kappa^+} : \text{there is no countable } \sigma \subseteq \kappa^{++} \text{ as above for } M_i\}$

cannot be stationary: Suppose  $S$  is stationary. Let  $\alpha \in \kappa^{++}$ ,  $\text{cf}(\alpha) = \omega$ , such that  $F''[\alpha]^{<\omega} \cap \kappa^{++} \subseteq \alpha$  for some  $F : [H(\lambda)]^{<\omega} \rightarrow H(\lambda)$  generating club of elementary submodels  $R$  such that  $(M_j)_{j < \kappa^+} \in R$ .

Now we can easily find  $X \subseteq \alpha$  cofinal in  $\alpha$ , such that  $R = F''[X]^{<\omega}$  is such that  $|R| = \kappa$  and  $i := R \cap \kappa^+ \in S$ . Let  $\sigma \subseteq X$  be countable and cofinal in  $X$ . But then  $R$  is a  $\kappa^+$ -end-extension of  $M_i$  and  $\text{cf}(R \cap \kappa^{++}) = \omega$ , and so  $\sigma$  witnesses that  $M_i \notin S$ . Contradiction.  $\square$

Now we get club-many  $i$  such that if  $\text{cf}(i) = \kappa$ , then  $\text{cf}(M_i \cap \kappa^{++}) = \omega$ . But this is impossible since  $(\sup(M_i \cap \kappa^{++})) : i < \kappa^+$  is strictly increasing and continuous and therefore  $\text{cf}(M_i \cap \kappa^{++}) = \kappa > \omega$  if  $\text{cf}(i) = \kappa$ .  $\square$

# Saturation

Given an infinite regular  $\kappa$  and a stationary  $S \subseteq \kappa^+$ ,  $\text{NS}_{\kappa^+} \upharpoonright S$  is *saturated* iff every collection  $\mathcal{A}$  of stationary subsets of  $S$  such that  $S_0 \cap S_1$  is nonstationary for all  $S_0 \neq S_1$  in  $\mathcal{A}$  is such that  $|\mathcal{A}| \leq \kappa^+$ .

## Fact

If  $\kappa$  is an infinite regular cardinal,  $SRP(\kappa^+, 1)$  implies that  $NS_{\kappa^+} \upharpoonright S_{\kappa^+}^{\kappa^+}$  is saturated.

**Proof:** Let  $\mathcal{A}$  be a collection of stationary subsets of  $S_{\kappa^+}^{\kappa^+}$  with pairwise nonstationary intersection. We want to show  $|\mathcal{A}| \leq \kappa^+$ . Let  $X = \mathcal{A} \cup \kappa^+$  and let  $\mathcal{S}$  be the collection of  $Z \in [X]^\kappa$  such that

- $\delta_Z := Z \cap \kappa^+ \in \kappa^+$  and
- $\delta_Z \in S$  for some  $S \in \mathcal{A} \cap Z$ .

Let  $(M_i)_{i < \kappa^+}$  be a reflecting sequence for  $\mathcal{S}$  as given by  $SRP(\kappa^+, 1)$ , and suppose  $S \in \mathcal{A} \setminus \bigcup_{i < \kappa^+} M_i$ . Let  $M'_i = \text{Hull}_\lambda(M_i \cup \{S\})$  for all  $i$  and note that

$$\{i < \kappa^+ : \text{cf}(i) = \kappa \Rightarrow M'_i \cap \kappa^+ = M_i \cap \kappa^+\}$$

contains a club  $C \subseteq \kappa^+$ .



Hence, for every  $i \in C \cap S$  there is some  $S(i) \in M_i$  such that  $M_i \cap \kappa^+ \in S(i)$ . By Fodor's lemma there is some  $S_0$  such that

$$T = \{i \in S \cap C : S(i) = S_0\}$$

is stationary. But that is a contradiction since  $M_i \cap \kappa^+ \in S \cap S_0$  for every  $i \in T$  and therefore  $S \cap S_0$  is stationary.  $\square$

Let us say that a forcing  $\mathcal{P}$  is  $\kappa$ -strongly 1-semiproper iff it satisfies the definition of ' $\kappa$ -strongly semiproper' replacing  $\text{Hull}(M, \sigma)$ , for  $|\sigma| \leq \kappa$ , with  $\text{Hull}(M, \sigma)$ , for  $|\sigma| \leq 1$ .

$\kappa$ -strong 1-semiproperness is the least demanding excursion of  $\kappa$ -strong properness into the realm of semiproperness.

$\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice, } \kappa\text{-strongly 1-semiproper}\})$

implies  $\text{SRP}(\kappa^+, 1)$  and therefore the saturation of  $\text{NS}_{\kappa^+} \upharpoonright \mathcal{S}_{\kappa}^{\kappa^+}$ .

**Question:** Is  $\text{FA}_{\kappa^+}(\{\mathcal{P} : \mathcal{P} \text{ } \kappa\text{-lattice, } \kappa\text{-strongly 1-semiproper}\})$  consistent for any  $\kappa \geq \omega_1$ ?

**Question:** Suppose  $\kappa \geq \omega_1$  is regular and  $\text{NS}_{\kappa^+} \upharpoonright \mathcal{S}_{\kappa}^{\kappa^+}$  is saturated. Does it follow that GCH cannot hold below  $\kappa$ ?

## On high properness when adding reals

Neeman considers side conditions consisting of *nodes* of either of the following types.

- (1) (Countable type elementary) These are models  $M \preceq H(\theta)$  such that  $|M| = \aleph_0$ .
- (2) (Type  $\omega_1$ ) These are IC models  $N \preceq H(\theta)$  such that  $|N| = \aleph_1$ .
- (3) (Countable type tower.) These are countable  $\in$ -chains  $\mathcal{T}$  of nodes of type  $\omega_1$  such that  $\mathcal{T} \cap N \in N$  for all  $N \in \mathcal{T}$ .

## Definition

(Neeman) A *two-size side condition* is a finite set  $\mathcal{N}$  of nodes of the above types which is  $\in$ -increasing (i.e., every node belongs to the next), and closed under intersection in the sense that:

- If  $N, M \in \mathcal{N}$ ,  $N \in M$ ,  $N$  of type  $\omega_1$ , and  $M$  countable elementary, then  $M \cap N \in \mathcal{N}$ .
- If  $N, \mathcal{T} \in \mathcal{N}$ ,  $N \in \mathcal{T}$ ,  $\mathcal{T}$  of type tower, and  $\mathcal{T} \cap N \neq \emptyset$ , then there is a tower  $\mathcal{T}' \supseteq \mathcal{T} \cap N$  occurring in  $\mathcal{N}$  before  $N$ .

## Definition

(Neeman) A partial order  $\mathcal{P}$  is *two-size proper* if for every large enough  $\theta$  there is a function  $f : [H(\theta)]^{<\omega} \rightarrow H(\theta)$  such that for every two-size side condition  $\mathcal{N}$  with all models involved closed under  $f$ , every  $Q \in \mathcal{N}$ , and every  $p \in \mathcal{P} \cap Q$ , if  $p$  is  $(R, \mathcal{P})$ -generic for every  $R \in \mathcal{N} \cap Q$ , then there is  $q \leq_{\mathcal{P}} p$  which is  $(R, \mathcal{P})$ -generic for all  $R \in \mathcal{N}$ . (If  $\mathcal{T}$  is a tower, a condition is  $(\mathcal{T}, \mathcal{P})$ -generic iff it is  $(N, \mathcal{P})$ -generic for all  $N \in \mathcal{T}$ .)

## Theorem

(Neeman) If  $\kappa$  is a supercompact cardinal, then there is a partial order  $\mathcal{P} \subseteq V_{\kappa}$  forcing  $FA_{\aleph_2}(\{\mathcal{P} : \mathcal{P} \text{ two-size proper}\})$ .

A partial order  $\mathcal{P}$  is *two-size strongly semiproper* if for every large enough  $\theta$  there is a function  $f : [H(\theta)]^{<\omega} \rightarrow H(\theta)$  such that for every two-size side condition  $\mathcal{N}$  with all models involved closed under  $f$ , every  $Q \in \mathcal{N}$ , and every  $p \in \mathcal{P} \cap Q$ , if  $p$  is  $(R, \mathcal{P})$ -strongly  $\omega_2$ -semigeneric for every  $R \in \mathcal{N} \cap Q$ , then there is  $q \leq_{\mathcal{P}} p$  which is  $(R, \mathcal{P})$ -strongly  $\omega_2$ -semigeneric for all  $R \in \mathcal{N}$ .

## Theorem

$FA_{\aleph_2}(\{\mathcal{P} : \mathcal{P} \text{ two-size strongly semiproper}\})$  implies  $SRP(\omega_2, \omega)$ .

## Corollary

$FA_{\aleph_2}(\{\mathcal{P} : \mathcal{P} \text{ two-size strongly semiproper}\})$  is inconsistent.

Two-size strong 1-semiproperness is the least demanding excursion of two-size properness into the realm of semiproperness.

$\text{FA}_{\aleph_2}(\{\mathcal{P} : \mathcal{P} \text{ two-size strongly 1-semiproper}\})$  implies  $\text{SRP}(\omega_2, 1)$ .

**Question:** Is  $\text{FA}_{\aleph_2}(\{\mathcal{P} : \mathcal{P} \text{ two-size strongly 1-semiproper}\})$  consistent?

In joint work with Veličković, and using forcing with virtual models with generators, we do get consistency of a shadow of  $\text{SRP}(\omega_2, 1)$  but which unfortunately doesn't seem to be enough to get saturation of  $\text{NS}_{\omega_2} \upharpoonright \mathcal{S}_{\omega_1}^{\omega_2}$ .

# On high stationary reflection

## Theorem

(Sakai)

- (1)  $WRP_{\omega_1} \upharpoonright IA_\omega$  implies  $2^{\aleph_0} \leq \aleph_3$ .
- (2) If  $\kappa$  is supercompact, then the  $\aleph_1$ -support iteration of length  $\kappa$  with mixed support for collapsing  $\alpha$  to  $\omega_2$  (for  $\alpha < \kappa$ ) with conditions of size  $\aleph_1$  while also adding Cohen reals forces  $WRP_{\omega_1} \upharpoonright IA_\omega + 2^{\aleph_0} = \aleph_3$ .



Some final questions:

**Question:** Is there any consistent high analogue  $R^*$  of any reflection principle  $R$  following from  $MM^{++}$  such that  $R^*$  implies  $2^{\aleph_0} = \aleph_3$ ?

**Question:** Is there any  $\Pi_2$  sentence  $\sigma$  such that the following holds?

- (1) ZFC proves that if  $H(\omega_3) \models \sigma$ , then  $2^{\aleph_0} = \aleph_3$ .
- (2) For some reasonable large cardinal axiom LC, ZFC+ LC proves that it is forcible that  $H(\omega_3) \models \sigma$ .

Related to the last question in the previous slide but for  $H(\omega_2)$ :

**Conjecture:**  $\text{BFA}(\{Q : Q \text{ } \omega\text{-proper}\})$  implies  $2^{\aleph_0} = \aleph_2$ .